

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 043 976 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
14.12.2005 Bulletin 2005/50

(51) Int Cl.7: **A61K 9/20**, A61K 9/22,
A61K 9/52, A61K 31/44,
A61K 31/41

(21) Application number: **98964631.0**

(86) International application number:
PCT/SE1998/002368

(22) Date of filing: **17.12.1998**

(87) International publication number:
WO 1999/032091 (01.07.1999 Gazette 1999/26)

(54) ORAL PHARMACEUTICAL EXTENDED RELEASE DOSAGE FORM

ORALE PHARMAZEUTISCHE DOSIERUNGSFORM MIT VERLÄNGERTER
WIRKSTOFFFREISETZUNG

FORME POSOLOGIQUE PHARMACEUTIQUE ORALE A LIBERATION PROLONGEE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
LT LV MK RO SI

(73) Proprietor: **AstraZeneca AB**
151 85 Södertälje (SE)

(30) Priority: **22.12.1997 SE 9704869**

(72) Inventors:
• **KAREHILL, Per-Gunnar**
S-431 83 Mölndal (SE)
• **LUNDBERG, Per, Johan**
S-431 83 Mölndal (SE)

(43) Date of publication of application:
18.10.2000 Bulletin 2000/42

(56) References cited:
WO-A1-97/02020 WO-A1-97/47285

EP 1 043 976 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionField of the invention

[0001] The present invention is related to new pharmaceutical dosage forms which comprise a proton pump inhibitor, i.e. a H⁺,K⁺-ATPase inhibitor. The new dosage forms are enteric coated formulations which provide an extended and continuous release of the H⁺,K⁺-ATPase inhibitor in the small and/or large intestines resulting in an extended blood plasma profile. The formulations comprise a hydrophilic or hydrophobic matrix resulting in an extended release of the H⁺,K⁺-ATPase inhibitor for a minimum of 2 and a maximum of 12 hours. Furthermore, the present invention refers to the manufacture of such extended release pharmaceutical formulations, and their use in medicine.

Background of the invention and prior art

[0002] Acid labile H⁺, K⁺-ATPase inhibitors also named as gastric proton pump inhibitors are for instance compounds known under the generic names omeprazole, lansoprazole, pantoprazole, rabeprazole and leminoprazole. Some of these compounds are disclosed in EP-A1-0005129, EP-A1-124495, WO 94/27988, EP-A1-174726, EP-A1-166287 and GB 2163747.

[0003] These pharmaceutical substances are useful for inhibiting gastric acid secretion in mammals including man by controlling gastric acid secretion at the final step of the acid secretory pathway and thus reduce basal and stimulated gastric acid secretion irrespective of stimulus. In a more general sense, they may be used for prevention and treatment of gastric-acid related diseases in mammals and man, including e.g. reflux oesophagitis, gastritis, duodenitis, gastric ulcer, duodenal ulcer and Zollinger-Ellison syndrom. Furthermore, they may be used for treatment of other gastrointestinal disorders where gastric acid inhibitory effect is desirable e.g. in patients on NSAID therapy, in patients with Non Ulcer Dyspepsia, and in patients with symptomatic gastro-oesophageal reflux disease (GORD). They may also be used in patients in intensive care situations, in patients with acute upper gastrointestinal bleeding, pre-and postoperatively to prevent aspiration of gastric acid and to prevent and treat stress ulceration. Further, they may be useful in the treatment of psoriasis as well as in the treatment of *Helicobacter* infections and diseases related to these.

[0004] Therapeutic control of gastric acid secretion is fundamental in all these diseases, but the degree and duration of acid inhibition required for optimal clinical effect is not fully understood.

[0005] It has been proposed by the Applicant in WO97/48380 (published 24 December, 1997, i.e. after the priority date of the present application) that an administration regimen that gives blood plasma levels extending from 2 to 12 hours (by any of several means) will result in a larger fraction of the proton pumps being inhibited. Thus, an extended blood plasma level should result in more effective inhibition of acid secretion resulting in improved efficacy in GORD, more rapid healing of gastric ulcer and improved eradication of *H. Pylori*. The present invention provides pharmaceutical dosage forms which achieve such extended plasma levels by an extended release of the drug.

[0006] A pharmaceutical dosage form comprising omeprazole or any other proton pump inhibitor is best protected from contact with acidic gastric juice by an enteric coating layer. In US 4,786,505 and US 4,853,230 such enteric coated preparations are described. These preparations have a core comprising an alkaline salt of the drug or a core comprising the drug together with an alkaline reacting compound, the core is coated with a water soluble or in water rapidly disintegrating separating layer and then with an enteric coating layer. WO 96/01623 and WO 96/01624 describe tableted dosage forms of omeprazole and other proton pump inhibitors, wherein enteric coating layered pellets are compressed into a multiple unit tableted dosage form. It is essential in these tableted formulations that the enteric coating layer can withstand the compression forces. None of these by the Applicant previously described formulations gave an extended release of the drug which resulted in an extended blood plasma profile.

[0007] WO 97/02020 describes a dosage form for pantoprazol together with an antibiotic substance, which dosage form has a release-slowing membrane positioned as a intermediate layer. Said membrane comprises a water-insoluble film-forming agent as an important feature of the dosage forms. WO 97/02021 describes the same type of dosage form for a reversible proton pump inhibitor in combination with an antibiotic substance.

[0008] A facilitated way to produce extended release dosage forms compared to applying a semipermeable membrane, is to make a dosage form comprising a matrix unit. Some advantages of such matrices are for instance easier processing methods mainly by the use of common granulating and tableting equipment, and sometimes also with regard to solvents handling, energy and production time gain etc.

[0009] The use of hydrophilic matrix tablets as a principle for extended drug release was first described in the early 60's, see for instance US Patent 3,065,143. Also the hydrophobic matrix tablet principle for extended release originates from the 60's, for instance quinidine dures were on the market in 1963.

[0010] Extended release dosage forms comprising different drugs in a matrix have been described in prior art. However, none of these matrix dosage forms as such is suitable for a H⁺,K⁺-ATPase inhibitor.

[0011] Some extended release hydrophilic matrix dosage forms are described in the literature for instance: In Journal

of Pharmaceutical Sciences vol. 84, No. 3, March 1995, in which Kim describes dosage forms comprising theophylline or diltiazem hydrochloride. US Patent 5,273,758 describes dosage forms comprising for instance clemastine fumarate. EP 0249587 discusses felodipine formulations. Dosage forms comprising a benzodiazepine derivative are described by Franz et al in Journal of Controlled Release 1987, 5, 159-72.

[0012] Dosage forms comprising an extended release hydrophobic matrix have been described for instance by Romero et al in International Journal of Pharmacy 1991, 73, 239-48.

[0013] Extended release tablets with an additional coating layer have also been described, for instance by Sangalli et al in International Journal of Pharmaceutics, 91(1993), 151-6. The drugs exemplified are metoprolol tartrate and benfluorex. The described dosage form has an impermeable coating which is perforated to achieve a hole in the middle of the tablet, exposing a starting surface area for the dissolution of the inner core, i.e. dissolution of the active drug.

[0014] A rather complicated dosage form was described in US Patent 5,178,867. The dosage forms had a core comprising a drug which core was coated with a semipermeable wall (maintaining its physical integrity during the life-time of the dosage form) having at least one hole drilled through it as an exit port for the dissolved drug. It is also mentioned that an enteric coating layer may be applied for restricting drug delivery in the stomach and for providing drug release in the small intestine. This dosage form is much more complicated to manufacture than a matrix unit. There is no detailed description of a prepared dosage form comprising a proton pump inhibitor compound and testing of such a dosage form to assure that no acidic gastric fluid is penetrating the semipermeable membrane, and that the active substance is delivered intact to the site of absorption.

[0015] None of these dosage forms provides an easy-to-produce matrix dosage form which protect an acidic susceptible substance such as a proton pump inhibitor against degradation which occurs in contact with an acidic milieu such as the one found in the stomach.

Summary of the invention

[0016] Thus, the present invention relates to an enteric coated formulation with extended release properties comprising a hydrophilic or hydrophobic matrix, in which a H^+, K^+ -ATPase inhibitor or one single enantiomer thereof, or an alkaline salt of the H^+, K^+ -ATPase inhibitor or one of its single enantiomers is incorporated.

[0017] The present invention provides a solution to the problem of making in a simplified manner such extended release dosage forms comprising an acidic susceptible H^+, K^+ -ATPase inhibitor, such as omeprazole or another proton pump inhibitor. A specific problem is that the pharmaceutical dosage forms according to the present invention must fulfill certain requirement with respect to gastric acid resistance for enteric coated articles specified in the US Pharmacopeia (Edition 23). Such as the dosage form has to be protected by an enteric coating to ensure safe delivery of the intact drug to the proper site in the gastrointestinal channel where it may be absorbed.

[0018] According to the present invention the extended plasma profile is provided by once daily administration of an enteric coated dosage form which releases the proton pump inhibitor during an extended time period, during a minimum period of 2 hours and a maximum period of 12 hours. Thus, the complete dose shall have been delivered within 2 hours or at a maximum within 12 hours. The therapeutic effect of omeprazole and similar substances may be improved by providing an extended plasma profile and by providing such a dosage form for a once daily administration.

[0019] The present extended release formulations show an improved patient compliance over an administration regimen comprising consecutive administration of two or more unit doses during one day.

Detailed description of the invention.

[0020] The dosage forms giving extended release according to the present invention, are units in the form of enteric coated tablets. Alternatively, the units are enteric coated pellets, which pellets are filled into a capsule or together with tablet excipients compressed into a multiple unit tableted dosage form.

[0021] The individual units, i.e. tablets or pellets, may be constructed as a

- a core material, optionally layered on a seed/sphere, the core material comprises a hydrophilic or hydrophobic matrix containing the active drug and optionally pharmaceutically acceptable excipients, and
- an optional surrounding separating layer, and finally
- an enteric coating layer.

Core material.

[0022] The core material for the units, i.e. the tablets or the individual pellets can be constituted according to different principles. The core material may be homogenous or heterogeneous.

I) *Homogenous core material.*

[0023] If the core material is homogenous, it has a homogenous distribution of active substance throughout the core material.

[0024] The active substance is mixed with substances forming a hydrophilic or hydrophobic matrix and optionally pharmaceutically acceptable excipients. The core material should be free from acidic substance. Thus, the hydrophilic or hydrophobic matrix in combination with other material in the core must not create an acidic reaction in the core material, which would be deleterious to the acid susceptible proton pump inhibitor compound. The micro environment around the proton pump inhibitor compound should preferably have a pH of not less than pH=7, more preferably not less than pH=8, when water is absorbed to the particles of the mixture or when water is added in small amount to the mixture.

[0025] The active substance may be mixed with further components to obtain preferred handling and processing properties and a suitable concentration of the active substance in the final mixture. Such components can be binders, surfactants, lubricants, glidants, fillers, alkaline additives or other pharmaceutically acceptable ingredients, alone or in mixtures.

[0026] Said core material may be produced either by direct compression of the mixed ingredients, or by granulation of the ingredients followed by compression of the dried granulated material.

[0027] In direct compression, the ingredients are mixed and compressed by using ordinary tableting equipment.

[0028] For the granulation there are numerous alternatives of granulating procedures mentioned in the literature, dry methods like roller compaction (Chilsonator) and wet methods utilizing granulating solutions with and without the addition of binders. A variant of the wet methods is to make a spray-granulation in a fluid bed.

[0029] For the wet granulating methods either organic solvents, aqueous solutions or pure water may be utilized to prepare the granulating solutions. Due to environmental considerations pure water is preferred. However, for some of the materials used as hydrophilic matrix components, the technical properties of the produced granules might be better when using organic solvents such as alcohols, this is especially noticeable for hydroxypropyl methylcelluloses.

[0030] For granulation of the hydrophobic matrix components it is also preferred to use alcoholic solvents in wet granulation methods. As binders in these solution, one or more of the polymers listed below, as matrix forming polymers may be chosen.

[0031] As a general principle the active ingredients together with matrix forming polymers and optionally pharmaceutically acceptable excipients are mixed and granulated. Dried granules are optionally mixed with pharmaceutically acceptable excipients, and then compressed to tablets utilizing common equipment.

[0032] The size of the formulated core materials is approximately between 2 and 14 mm, preferably between 3 and 9 mm for a tablet preparation, and between 0.1 and 4 mm, preferably between 0.1 and 2 mm for a pellet preparation.

II) *Heterogenous core material.*

[0033] Alternatively, the core material may be heterogeneous with an inner zone, for instance a seed or sphere, not containing the active substance. This seed or sphere is surrounded by a layer of a hydrophilic or hydrophobic matrix containing the active substance, and optionally pharmaceutically acceptable excipients are incorporated in the matrix.

[0034] The seed or sphere may be soluble or insoluble. Optionally, the seed or sphere (inner zone) may be coated with an inert layer to prepare a smooth surface before the layer containing active substance and hydrophilic or hydrophobic eroding substance(s) is applied onto the seed/sphere.

[0035] Insoluble seeds/spheres may comprise different oxides, celluloses, organic polymers and other materials, alone or in mixtures. Water soluble seeds/spheres may comprise different inorganic salts, sugars and other materials, alone or in mixtures. The size of the seeds may vary between approximately 0.1 and 2 mm. The seeds layered with the matrix containing the active substance are produced either by powder or solution/suspension layering using for instance granulating or spray coating/layering equipment.

Pharmaceutically acceptable additives.

[0036] Binders for a hydrophilic matrix can be chosen among the hydrophilic eroding matrices mentioned below, and in addition from sugars, polyvinyl pyrrolidone, starches and gelatine.

[0037] Binders for a hydrophobic matrix can be chosen among the hydrophobic eroding matrices mentioned below.

[0038] Additives listed among the following components are suitable both for a hydrophilic as well as a hydrophobic matrix.

[0039] Suitable alkaline additives can be chosen among, but are not restricted to, substances such as the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric acid, carbonic acid, citric acid or other suitable weak inorganic or organic acids; aluminium hydroxide/sodium bicarbonate coprecipitate; substances normally used in

antacid preparations such as aluminium, calcium and magnesium hydroxides; magnesium oxide or composite substances, such as $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$, $(\text{Mg}_6\text{Al}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O})$, $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ or similar compounds; organic pH-buffering substances such as trihydroxymethylaminomethane, basic amino acids such as arginine, and their salts or other similar pharmaceutically acceptable pH-buffering substances.

[0040] Suitable surfactants are found in the groups of pharmaceutically acceptable non-ionic surfactants, such as polysorbate 80, or ionic surfactants such as for instance sodium lauryl sulfate.

[0041] Lubricants are for instance magnesium stearate, sodium stearyl fumarate (Pruv™), and cetyl palmitate.


[0042] Fillers are for instance sodium aluminium silicate, lactose, calcium phosphate, and others.

[0043] Glidants are for instance talc and aerosil.


[0044] Antioxidants may be added when appropriate.

Active substance.


[0045] Compounds of interest for the novel extended release dosage forms according to the present invention are compounds of the general formula I, an alkaline salt thereof, one of the single enantiomers thereof or an alkaline salt of one of the enantiomers




wherein
Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R_6 - R_9 optionally may be exchanged for a nitrogen atom without any substituents;

R_1 , R_2 and R_3 are the same or different and selected from hydrogen, alkyl, alkoxy optionally substituted by fluorine, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R_4 and R_5 are the same or different and selected from hydrogen, alkyl and arylalkyl;




R_6 is hydrogen, halogen, trifluoromethyl, alkyl or alkoxy;





R_6 - R_9 are the same or different and selected from hydrogen, alkyl, alkoxy, halogen, haloalkoxy, alkylcarbonyl, alkoxycarbonyl, oxazolinyl, trifluoroalkyl, or adjacent groups R_6 - R_9 form ring structures which may be further substituted;



R_{10} is hydrogen or forms an alkylene chain together with R_3 and

R_{11} and R_{12} are the same or different and selected from hydrogen, halogen or alkyl.

[0046] Examples of specifically interesting compounds according to formula I are







[0047] The compound suitable to be used the extended release formulations according to the present invention may be used in neutral form or in the form of an alkaline salt, such as for instance the Mg^{2+} , Ca^{2+} , Na^{+} or K^{+} salts, preferably the Mg^{2+} salts. The compounds may also be used in the form of one of its single enantiomers or an alkaline salt of the single enantiomer.

[0048] Preferred compounds for the oral pharmaceutical preparations according to the present invention are omeprazole, a magnesium salt of omeprazole or a magnesium salt of the (-)-enantiomer of omeprazole. Omeprazole and related substances as well as their preparations are described in EP 5129, EP 124 495, WO 95/01977, WO 94/27988 hereby incorporated in a whole by references.

[0049] The above compounds are susceptible to degradation/transformation in acidic and neutral media. Generally, the degradation is catalyzed by acidic reacting compounds and the active compounds are stabilized with alkaline reacting compounds. There are different enteric coating layered preparations comprising omeprazole as well as other proton pump inhibitors described in the prior art, see for instance US-A 4,853,230, WO 95/01783 and WO 96/01623. Especially, the latter describes alternative manufacturing methods for the preparation of enteric coating layered pellets comprising omeprazole and similar compounds. These patents are hereby incorporated in whole by references.

Hydrophilic matrix.

[0050] The active substance, i.e. the drug, is embedded in a hydrophilic polymer optionally together with pharmaceutically acceptable excipients. Suitable hydrophilic polymers are for instance hydroxypropyl methylcellulose, hydroxypropyl cellulose, ethylhydroxy ethylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, polyethylene oxides, polyvinyl alcohols, tragacanth, and xanthan. These polymers can be used alone or in mixtures with each other.

[0051] The amount of hydrophilic polymer in the matrix is preferably 15-80 % (calculated on the unit weight), and the hydrophilic polymer(s) chosen among the above mentioned.

[0052] Especially preferred polymers in the hydrophilic matrix unit are hydroxypropyl methylcellulose or polyethylene oxides.

[0053] Excipients preferred in the matrix are fillers which result in good technical tableting properties, i.e. sodium aluminium silicate, mannitol or calcium phosphate (Emcompress™). A preferred matrix comprises 15-80% w/w (calculated on the unit weight) of a hydrophilic polymer chosen as above, and 10-60% w/w (calculated on the unit weight) of sodium aluminium silicate or calciumphosphate (Emcompress™).

Hydrophobic matrix.

[0054] The active substance, i.e. the drug, is embedded in a hydrophobic matrix optionally together with pharmaceutically acceptable excipients. The hydrophobic matrix comprises a hydrophobizing agent and/or a hydrophobic polymer. Suitable material for the hydrophobic matrix are for instance a hydrophobizing agents such as cetanol, ceto-stearyl alcohol, cetyl palmitate, waxes like carnauba wax, paraffin, magnesium stearate, sodium stearyl fumarate, and medium- or long- chain glycerol esters alone or in any mixtures. Hydrophobic polymers are exemplified by for instance polyvinyl chloride, ethyl cellulose, polyvinyl acetate and acrylic acid copolymers, such as Eudragith™ RS and RL. The polymers can be used alone or as mixtures.

[0055] As binders for the hydrophobic matrix may be used either hydrophilic or hydrophobic polymers.

[0056] It is important that the matrix comprises at least one component that is soluble in media such as the intestinal fluids. This component dissolves and leaves a porous network open for passage of dissolving fluids and dissolved drug. This soluble component may be the active drug itself, or a soluble component such as a sugar. Preferably the soluble component is present in an amount of not less than 2% w/w (calculated on the unit weight) and up to 60%.

[0057] It is preferred that the matrix comprises not less than 10 % w/w (calculated on the unit weight) and up to 80% of a hydrophobizing agent or a hydrophobic polymer, both described above, or any combinations thereof.

[0058] Another preferred matrix comprises as an additive a slightly soluble or less soluble component. As such components may any of the following be added: sodium aluminium silicate, calciumphosphate, aerosil, titaniumdioxide, magnesium carbonates, or other neutral or alkaline compounds that are slightly soluble or less soluble, herein with regard to solubility in water. Slightly soluble is defined in compliance with the European Pharmacopoeia (Edition 3) under the heading "General notices". Such a matrix comprises 10-80 % w/w (calculated on the unit weight) of a hydrophobizing agent or a hydrophobic polymer or any combinations thereof, together with 10% - 60% of a slightly soluble or less soluble component. As such a component is especially preferred sodium aluminium silicate.

[0059] The final dissolution profile may sometimes be adjusted by thermal treatment of the hydrophobic matrix unit for a short period, to achieve temperatures at or above the softening temperature of the hydrophobizing agents. Such a treatment is most suitably performed after the enteric coating has been completed.

Enteric coating layer(s) and separating layer(s).

[0060] Before applying an enteric coating layer onto the core material, the pellet or tablet may optionally be covered with one or more separating layers comprising pharmaceutical excipients optionally including alkaline compounds such as for instance pH-buffering compounds. This separating layer separates the active substance in the pellets or tablets from the outer enteric coating layer.

[0061] The separating layer can be applied to by coating or layering procedures in suitable equipments such as coating pan, coating granulator, centrifugal granulator in a fluidized bed apparatus (including Wurster type) using water and/or organic solvents for the coating process. As an alternative the layer(s) can be applied by using powder coating or press-coating techniques.

[0062] Suitable materials for the separating layer are pharmaceutically acceptable compounds such as, for instance, sugar, polyethylene glycol, polyvinyl pyrrolidone, polyvinyl alcohol, polyvinyl acetate, hydroxypropyl cellulose, methylcellulose, ethylcellulose, hydroxypropyl methylcellulose, carboxymethylcellulose sodium and others, used alone or in mixtures. Additives such as plasticizers, colorants, pigments, fillers, anti-tacking and anti-static agents, such as for instance magnesium stearate, titanium dioxide, talc, pH-buffering substances and other additives may also be included into the separating layer.

[0063] When the optional separating layer is applied to the pellets or tablets it may constitute a variable thickness. The maximum thickness of the optional separating layer is normally only limited by processing conditions. The separating layer may serve as a diffusion barrier and may act as a pH-buffering zone. The optionally separating layer may improve the chemical stability of the active substance and/or the physical properties of the dosage form.

[0064] Finally the units, i.e. the tablets or pellets, are covered by one or more enteric coating layers by using a suitable coating technique. The enteric coating layer material may be dispersed or dissolved in either water or in suitable organic solvents. As enteric coating layer polymers one or more, separately or in combination, of the following can be used; e.g. solutions or dispersions of methacrylic acid copolymers, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, cellulose acetate trimellitate, carboxymethyl ethylcellulose, shellac or other suitable enteric coating layer polymer(s).

[0065] Additives such as dispersants, colorants, pigments, additional polymers e.g. poly(ethylacrylat, methylmethacrylat), anti-tacking and anti-foaming agents may also be included into the enteric coating layer. Other compounds may be added to increase film thickness and to decrease diffusion of acidic gastric juices into the acid susceptible material. The enteric coating layer(s) constitutes a thickness of approximately at least 10 µm, preferably more than 20 µm. The maximum thickness of the applied enteric coating layer(s) is normally only limited by processing conditions.

[0066] The enteric coating layers may also contain pharmaceutically acceptable plasticizers to obtain desired mechanical properties. Such plasticizers are for instance, but not restricted to, triacetin, citric acid esters, phthalic acid esters, dibutyl sebacate, cetyl alcohol, polyethylene glycols, glucerol monoesters, polysorbates or other plasticizers and mixtures thereof. The amount of plasticizer is preferably optimized for each formula, in relation to the selected polymer(s), selected plasticizer(s) and the applied amount of said polymer(s).

Final dosage form

[0067] The enteric coated tablet, or pellet optionally mixed with tablet excipients are filled into a capsule, or compressed into a multiple unit tableted dosage form. Prepared enteric coated tablets are optionally covered with filmforming agent(s) to obtain a smooth surface of the tablet and further enhance the stability of the tablet during packaging and transport. Such a tablet coating layer may further comprise additives like anti-tacking agents, colorants and pigments or other additives to obtain a tablet of good appearance.

[0068] The dosage forms according to the invention are suitable for oral administration. The dose will depend on the nature and severity of the disease to be treated. The dose may also vary according to the age, body weight, and response of the individual patient. Children and patients with liver diseases as well as patients under long term treatment will generally benefit from doses that are somewhat lower than the average. In the treatment of other conditions higher doses than average will be used. The dosage form may also be used in combinations with other dosage forms comprising for instance NSAID(s), motility agents, antibacterial substances, and/or antacids.

[0069] A unit dosage of the proton pump inhibitor is administered at least once a day. The oral pharmaceutical formulation will maintain an extended release of the pharmaceutical substance of a minimum of 2 and a maximum of 12 hours, preferably is maintained for a minimum of 4 and a maximum of 8 hours. Such an extended release preparation may comprise up to 500 mg of the substance, preferably the doses comprise about 5 - 100 mg of the substance, and more preferably 10 - 80 mg.

Examples

[0070] The invention is described more in detail by the following non-limiting examples.

Example 1.

[0071] Extended release matrix tablets comprising Omeprazole -Mg (approx. 20 mg).

[0072] Granules for tablet cores were made according to the following composition (parts by weight);

Omeprazole-Mg	45
Polyethylene oxide (mwt approx. 4000 000), Polyox® WSR 301	195
ethanol 95% (w/v)	97

[0073] The powders were mixed in a mixer after which the ethanol was added in an even stream. The mass was dried in a drying oven at 50°C.

[0074] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight) ;

Granules for tablet core	235
Sodium stearyl fumarate (Pruv®)	1

[0075] The mixing was performed in a Kenwood mixer, and the mixture was compressed to tablets (6 mm in diameter) having an average weight of 123 mg, on a single punch tableting machine (Diaf).

The dissolution rate was tested by analyzing individual tablets using USP dissolution apparatus No. 2 (paddle) equipped with a stationary basket and operated at 100 rpm and 37°C. The dissolution medium was phosphate buffer pH 6.8.

The release rate obtained (n=2) is shown in table below;

<u>Time (Hours)</u>	<u>Released (% of dose)</u>
0.5	4-4
1	7-8

(continued)

Time (Hours)	Released (% of dose)
3	20-21
5	31-33
10	59-67
15	84-86
20	95-96

[0076] The prepared tablets can be further processed according to Example 3 or 4, i.e. apply an enteric coating on the tablet.

Example 2.

[0077] Extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 32 mg).

[0078] Granules for tablet cores were made according to the following composition (parts by weight);

S-omeprazole Mg-salt	300
Hydroxypropyl methylcellulose 50 cps	80
ethanol 95% (w/v)	356
Polyvinyl pyrrolidone K-90	40

[0079] The powders were mixed in a mixer after which the ethanol was added in an even stream. The mass was dried in a drying oven at 50°C.

[0080] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight);

Granules for tablet core	380
Sodium stearyl fumarate (Pruv®)	4

[0081] The mixing was performed in a Kenwood mixer whereafter the mixture was compressed to tablets (7 mm in diameter) having an average weight of 175 mg, on a single punch tableting machine (Diaf).

[0082] The prepared tablets can be further processed according to Example 3 or 4, i.e. apply an enteric coating on the tablet.

Example 3.

[0083] Enteric coated extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 32 mg).

[0084] Tablets from example 2 were coated first with a separating layer in a fluidized bed coating apparatus with a coating suspension of the following composition;

EtOH 99.5% (w/v)	85 parts by weight
Water purified	85 parts by weight
Hydroxypropyl methylcellulose 6 cps	10 parts by weight
Talc, micronized	2 parts by weight
Sum:	182 parts.

[0085] 200 grams of tablets were processed and the coating was continued until average tablet weight was 181 mg.

[0086] The tablets coated with a separating layer were coated with an enteric coating layer in the same equipment as for the preceding coating step. The coating solution used had the following composition;

Hydroxypropyl methylcellulose phthalate (HP-55®)	19 parts by weight
Cetanol	1 parts by weight

(continued)

Acetone	182 parts by weight
Ethanol (95% w/v)	78 parts by weight
Sum:	280 parts

[0087] 100 grams of the separating layer coated tablets were processed and the coating was continued until average tablet weight was 194 mg.

[0088] The tablets were exposed for 0.1 M HCl for 2 hours. The acid resistance was determined to 98%.

Example 4.

[0089] Enteric coated extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 32 mg).

[0090] The tablets obtained from Example 2 were directly coated with an enteric coating layer in a fluidized bed coating apparatus. The coating solution used had the following composition;

Hydroxypropyl methylcellulose phtalate (HP-55®)	19 parts by weight
Cetanol	1 parts by weight
Acetone	182 parts by weight
Ethanol (95% w/v)	78 parts by weight
Sum:	280 parts

[0091] 100 grams of the tablets were processed and the coating was continued until average tablet weight was 187 mg.

[0092] The tablets were exposed for 0.1 M HCl for 2 hours. The acid resistance was determined to 99%.

Example 5.

[0093] Extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 45 mg).

[0094] Granules for tablet cores were made according to the following composition (parts by weight);

S-omeprazole Mg-salt	45
Polyethylene oxide (mwt approx. 4000 000), Polyox® WSR 301	145
Sodium aluminium silicate	50
Propyl gallate	0.1
Ethanol 99.5% (w/v)	140

[0095] The powders were mixed and moistened with the ethanol in a mixer after which the mass was dried in a drying oven at 50°C.

[0096] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight);

Granules for tablet core	232
Sodium stearyl fumarate (Pruv®)	1

[0097] The ingredients were mixed whereafter the mixture was compressed to tablets (10 mm in diameter) having an average weight of 241 mg, on a single punch tableting machine (Diaf).

[0098] Dissolution rate was tested as described in example 1. The release rate obtained (n=2) is shown in table below;

Time (Hours)	Released (% of dose)
2	16-16
4	29-29
6	41-42

(continued)

Time (Hours)	Released (% of dose)
8	53-54
10	65-66
12	76-78
14	88-88
16	95-96
18	100-100
20	109-109*)

*) Remark: the complete dose has been released.

Example 6.

[0099] Extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 45 mg).

[0100] Granules for tablet cores were made according to the following composition (parts by weight);

S-omeprazole Mg-salt	45
Polyethylene oxide (mwt approx. 4000 000), Polyox® WSR 301	72.5
Polyethylene oxide (mwt approx. 100 000), Polyox® WSR N10	72.5
Sodium aluminium silicate	50
Propyl gallate	0.1
Ethanol 99.5% (w/v)	140

[0101] The powders were mixed and moistened with the ethanol in a mixer after which the mass was dried in a drying oven at 50°C.

[0102] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight);

Granules for tablet core	234
Sodium stearyl fumarate (Pruv®)	1

[0103] The ingredients were mixed whereafter the mixture was compressed to tablets (10 mm in diameter) having an average weight of 241 mg, on a single punch tableting machine (Diaf).

[0104] Dissolution rate was tested as described in Example 1 above.

[0105] The release rate obtained (n=2) is shown in table below;

Time (Hours)	Released (% of dose)
2	14-14
4	29-29
6	44-47
8	60-65
10	73-78
12	87-89
14	99-101
16	101-102 *)
18	101-105 *)

*) Remark: The complete dose has been released

Example 7.

[0106] Extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 45 mg).

[0107] Granules for tablet cores were made according to the following composition (parts by weight);

S-omeprazole Mg-salt	45
Polyethylene oxide (mwt approx. 100 000), Polyox® WSR N10	145
Sodium aluminium silicate	50
Propyl gallate	0.1
Ethanol 99.5% (w/v)	140

[0108] The powders were mixed and moistened with the ethanol in a mixer after which the mass was dried in a drying oven at 50°C.

[0109] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following recipe (parts by weight);

Granules for tablet core	229
Sodium stearyl fumarate (Pruv®)	1

[0110] The ingredients were mixed whereafter the mixture was compressed to tablets (10 mm in diameter) having an average weight of 241 mg, on a single punch tableting machine (Diaf).

[0111] Dissolution rate was tested as described in example 1.

The release rate obtained (n=2) is shown in table below;

Time (Hours)	Released (% of dose)
2	67-68
4	107-110 *)
6	107-111 *)

*) Remark: The complete dose has been released.

Example 8.

[0112] Extended release matrix tablets comprising omeprazole Mg-salt (approx. 45 mg).

[0113] Granules for tablet cores were made according to the following composition (parts by weight);

Omeprazole Mg-salt	80
Hydroxypropyl methylcellulose 50 cps	300
Polyvinyl pyrrolidone K-90	40
Ethanol 99.5% (w/v)	400

[0114] The polyvinyl pyrrolidone (PVP) was dissolved in the alcohol. The other two ingredients were mixed and then moistened with the PVP-solution in a mixer. Thereafter the obtained mass was dried in a drying oven at 50°C.

[0115] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight);

Granules for tablet core	412
Sodium stearyl fumarate (Pruv®)	4

[0116] The ingredients were mixed whereafter the mixture was compressed to tablets (9 mm in diameter) having an average weight of 265 mg, on a single punch tableting machine (Diaf).

Example 9.

[0117] Extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 45 mg).

[0118] Granules for tablet cores were made according to the following composition (parts by weight);

S-omeprazole Mg-salt	74
Hydroxypropyl methylcellulose 50 cps	210
Hydroxypropyl methylcellulose 10000 cps	90
Polyvinyl pyrrolidone K-90	40
Ethanol 99.5% (w/v)	400

[0119] The polyvinyl pyrrolidone (PVP) was dissolved in the alcohol. The other ingredients were mixed and then moistened with the PVP-solution in a mixer. Thereafter the obtained mass was dried in a drying oven at 50°C.

[0120] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight);

Granules for tablet core	378
Sodium stearyl fumarate (Pruv®)	4

[0121] The mixing was performed in a mixer, and the mixture was compressed to tablets (9 mm in diameter) having an average weight of 261 mg, on a single punch tableting machine (Diaf).

[0122] Dissolution rate was tested in phosphate buffer pH 6.8 as described in example 1. The release rate obtained (n=6) is shown in table below;

Time (Hours)	Average (min-max) Released (% of nominal dose)
1	8 (8-8)
2	16 (16-17)
3	26 (25-27)
4	35 (34-36)
6	54 (52-56)
8	72 (70-75)
10	86 (83-91)
12	92 (90-99)

Example 10.

[0123] Extended release matrix tablets comprising S-omeprazole Mg-salt (approx. 55 mg).

[0124] Granules for tablet cores were made according to the following composition (parts by weight);

S-omeprazole Mg-salt	40
Polyvinyl alcohol mwt 22000, degree of hydrolysis 97.5-99.5%	160
Polyvinyl pyrrolidone K-90	14
Ethanol 99.5% (w/v)	49

[0125] The polyvinyl pyrrolidone (PVP) was dissolved in the alcohol. The other two ingredients were mixed and then moistened with the PVP-solution in a mixer. Thereafter the obtained mass was dried in a drying oven at 50°C.

[0126] After milling in an oscillating mill through a 1.0 mm screen the obtained granules were mixed with tablet lubricant, according to the following composition (parts by weight);

Granules for tablet core	215
Sodium stearyl fumarate (Pruv®)	2


[0127] The ingredients were mixed whereafter the mixture was compressed to tablets (9 mm in diameter) having an average weight of 310 mg, on a single punch tableting machine (Diaf).

[0128] Dissolution rate was tested in phosphate buffer pH 6.8 as described in example 1.


The release rate obtained (n=2) is shown in table below;

Claims


1. An enteric coated pharmaceutical extended release dosage form of a H⁺, K⁺-ATPase inhibitor, **characterized in that** the dosage form comprises a core material of a hydrophilic matrix or a hydrophobic matrix resulting in an extended release of the H⁺, K⁺-ATPase inhibitor for a minimum of 2 and a maximum of 12 hours and the H⁺, K⁺-ATPase inhibitor, and that the H⁺, K⁺-ATPase inhibitor is a compound of formula I, an alkaline salt thereof, one of the single enantiomers thereof or an alkaline salt of one of the enantiomers of a compound of formula I




wherein
Het₁ is




or




Het₂ is




or



X=



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from hydrogen, alkyl, alkoxy optionally substituted by fluorine,

alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₄ and R₅ are the same or different and selected from hydrogen, alkyl and arylalkyl;

R₆ is hydrogen, halogen, trifluoromethyl, alkyl or alkoxy;

R₆-R₉ are the same or different and selected from hydrogen, alkyl, alkoxy, halogen, haloalkoxy, alkylcarbonyl, alkoxy carbonyl, oxazoliny, trifluoroalkyl, or adjacent groups R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and

R₁₁ and R₁₂ are the same or different and selected from hydrogen, halogen or alkyl.

2. A dosage form according to claim 1 **characterized in that** the H⁺, K⁺-ATPase inhibitor is a compound selected from the group of omeprazole, an alkaline salt of omeprazole, the (-)-enantiomer of omeprazole and an alkaline salt of the (-)-enantiomer of omeprazole.

3. A dosage form according to claim 2, **characterized in that** the alkaline salt is a magnesium salt.

4. A dosage form according to claim 1 **characterized in that** the H⁺, K⁺-ATPase inhibitor is a compound selected from the group of lansoprazole, pantoprazole, alkaline salts thereof, a single enantiomer thereof, and an alkaline salt thereof.

5. A dosage form according to any of claims 1-4 **characterized in that** the core material is layered with a separating layer, which is present under the enteric coating layer.

6. A dosage form according to any of claims 1-5 **characterized in that** the core material further comprises pharmaceutically acceptable excipients.

7. A dosage form according to any of claims 1-6, **characterized in that** the core material further comprises alkaline additives.

8. A dosage form according to any of claims 1-7 **characterized in that** the core material comprises a seed layered with the H⁺, K⁺-ATPase inhibitor and the hydrophilic or hydrophobic matrix, and optionally pharmaceutically acceptable excipient.

9. A dosage form according to any of claims 1-8 **characterized in that** the core material creates a micro-environment around the H⁺, K⁺-ATPase inhibitor of not less than pH=7.

10. A dosage form according to any of claims 1-9 **characterized in that** the hydrophilic matrix comprises a hydrophilic polymer selected from the group of : hydroxypropyl methylcellulose, hydroxypropyl cellulose, ethylhydroxy ethylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, polyethylene oxides, polyvinylpyrrolidone, polyvinyl alcohols, tragacanth, and xanthan, or any mixtures thereof.

11. A dosage form according to claim 10, **characterized in that** the hydrophilic polymer is hydroxypropyl methylcellulose.

12. A dosage form according to claim 10 or 11 **characterized in that** the hydrophilic matrix further comprises a filler, such as sodium aluminium silicate.

13. A dosage form according to any of claims 1-9 **characterized in that** the hydrophobic matrix comprises a hydrophobic polymer and a hydrophobizing agent, the hydrophobizing agent is selected from the group of : cetanol, cetostearyl alcohol, cetyl palmitate, waxes like carnauba wax, paraffin, magnesium stearate, sodium stearyl fumarate, and medium- or long- chain glycerol esters alone or mixtures thereof.

14. A dosage form according to claim 13 **characterized in that** the hydrophobic polymer is selected from the group of : polyvinyl chloride, ethyl cellulose, polyvinyl acetate and acrylic acid copolymers.

15. A dosage form according to claims 13 or 14 **characterized in that** the hydrophobic matrix further comprises a slightly soluble or less soluble component selected from the group of : sodium aluminium silicate, calcium phosphate, aerosil, titanium dioxide, and magnesium carbonates.

16. A dosage form according to any of claims 1 - 15 **characterized in that** the H⁺, K⁺-ATPase inhibitor is a magnesium salt of the (-)-enantiomer of omeprazole.

17. A dosage form according to any of claims 1 - 16 for use in therapy.

18. A process for the manufacture of an enteric coated dosage form according to claim 1 comprising a H⁺, K⁺-ATPase inhibitor and a hydrophilic matrix or a hydrophobic matrix resulting in an extended release of the H⁺, K⁺-ATPase inhibitor for a minimum of 2 and a maximum of 12 hours, and optionally pharmaceutically acceptable excipient, **characterized in that** the process comprises the follow steps:

a) a core material is shaped comprising the H⁺, K⁺-ATPase inhibitor and the hydrophilic matrix or the hydrophobic matrix, and optionally pharmaceutically acceptable excipient,

b) optionally a separating layer is applied onto the core material, and


c) an enteric coating layer is applied onto the core material from step a) or step b).

19. Use of an oral pharmaceutical composition as claimed in any of claims 1 - 17 in the manufacture of a medicament with improved inhibition of gastric acid secretion.


20. Use of an oral pharmaceutical composition as claimed in any of claims 1 - 17 in the manufacture of a medicament with improved therapeutic effect in the treatment of gastrointestinal disorders associated with excess acid secretion.

Patentsprüche


1. Magensaftresistent beschichtete pharmazeutische Dosisform eines H⁺,K⁺-ATPase-Inhibitors mit verlängerter Wirkstofffreigabe, **dadurch gekennzeichnet, daß** die Dosisform ein Kernmaterial aus einer hydrophilen Matrix oder einer hydrophoben Matrix, die zu einer verlängerten Freigabe des H⁺,K⁺-ATPase-Inhibitors für mindestens 2 und höchstens 12 Stunden führt, und den H⁺,K⁺-ATPase-Inhibitor enthält und es sich bei dem H⁺,K⁺-ATPase-Inhibitor um eine Verbindung der Formel I, ein alkalisches Salz davon, eines der Einzelenantiomere davon oder ein alkalisches Salz eines der Enantiomere einer Verbindung der Formel I




worin
Het₁ für



steht,
Het₂ für



steht,
X =



worin

N in der Bezimidazolgruppierung bedeutet, daß eines der durch R₆-R₉ substituierten Ringkohlenstoffatome gegebenenfalls durch ein Stickstoffatom ohne Substituenten ersetzt sein kann;

R₁, R₂ und R₃ gleich oder verschieden sind und aus der Gruppe bestehend aus Wasserstoff, Alkyl, gegebenenfalls durch Fluor substituiertes Alkoxy, Alkylthio, Alkoxyalkoxy, Dialkylamino, Piperidino, Morpholino, Halogen, Phenyl und Phenylalkoxy ausgewählt sind;

R₄ und R₅ gleich oder verschieden sind und aus der Gruppe bestehend aus Wasserstoff, Alkyl und Arylalkyl ausgewählt sind;

R₆' für Wasserstoff, Halogen, Trifluormethyl, Alkyl oder Alkoxy steht;

R₆-R₉ gleich oder verschieden sind und aus der Gruppe bestehend aus Wasserstoff, Alkyl, Alkoxy, Halogen, Halogenalkoxy, Alkylcarbonyl, Alkoxycarbonyl, Oxazoliny und Trifluoralkyl ausgewählt sind oder benachbarte Gruppen R₆-R₉ gegebenenfalls weiter substituierte Ringstrukturen bilden;

R₁₀ für Wasserstoff steht oder gemeinsam mit R₃ eine Alkylkette bildet und

R₁₁ und R₁₂ gleich oder verschieden sind und aus der Gruppe bestehend aus Wasserstoff, Halogen oder Alkyl ausgewählt sind;

handelt.

2. Dosisform nach Anspruch 1, **dadurch gekennzeichnet, daß** es sich bei dem H⁺,K⁺-ATPase-Inhibitor um eine Verbindung aus der Gruppe bestehend aus Omeprazol, einem alkalischen Salz von Omeprazol, dem (-)-Enantiomer von Omeprazol und einem alkalischen Salz des (-)-Enantiomers von Omeprazol handelt.
3. Dosisform nach Anspruch 2, **dadurch gekennzeichnet, daß** es sich bei dem alkalischen Salz um ein Magnesiumsalz handelt.
4. Dosisform nach Anspruch 1, **dadurch gekennzeichnet, daß** es sich bei dem H⁺,K⁺-ATPase-Inhibitor um eine Verbindung aus der Gruppe bestehend aus Lansoprazol, Pantoprazol, alkalischen Salzen davon, einem Einzelantagonist davon und einem alkalischen Salz davon handelt.
5. Dosisform nach einem der Ansprüche 1-4, **dadurch gekennzeichnet, daß** das Kernmaterial mit einer unter der magensaftresistenten Schicht liegenden Trennschicht beschichtet ist.
6. Dosisform nach einem der Ansprüche 1-5, **dadurch gekennzeichnet, daß** das Kernmaterial ferner pharmazeutisch annehmbare Hilfsstoffe enthält.
7. Dosisform nach einem der Ansprüche 1-6, **dadurch gekennzeichnet, daß** das Kernmaterial ferner alkalische Additive enthält.

8. Dosisform nach einem der Ansprüche 1-7, **dadurch gekennzeichnet, daß** das Kernmaterial einen Keim umfaßt, der mit dem H⁺,K⁺-ATPase-Inhibitor und der hydrophilen oder hydrophoben Matrix und gegebenenfalls pharmazeutisch annehmbarem Hilfsstoff beschichtet ist.

9. Dosisform nach einem der Ansprüche 1-8, **dadurch gekennzeichnet, daß** das Kernmaterial eine Mikroumgebung um den H⁺,K⁺-ATPase-Inhibitor mit einem pH-Wert von mindestens 7 erzeugt.

10. Dosisform nach einem der Ansprüche 1-9, **dadurch gekennzeichnet, daß** die hydrophile Matrix zumindest teilweise aus einem hydrophilen Polymer aus der Gruppe bestehend aus Hydroxypropylmethylcellulose, Hydroxypropylcellulose, Ethylhydroxyethylcellulose, Hydroxyethylcellulose, Carboxymethylcellulose, Natriumcarboxymethylcellulose, Methylcellulose, Polyethylenoxiden, Polyvinylpyrrolidon, Polyvinylalkoholen, Tragant und Xanthan oder Gemischen davon besteht.

11. Dosisform nach Anspruch 10, **dadurch gekennzeichnet, daß** es sich bei dem hydrophilen Polymer um Hydroxypropylmethylcellulose handelt.

12. Dosisform nach Anspruch 10 oder 11, **dadurch gekennzeichnet, daß** die hydrophile Matrix ferner einen Füllstoff wie Natriumaluminiumsilicat enthält.

13. Dosisform nach einem der Ansprüche 1-9, **dadurch gekennzeichnet, daß** die hydrophobe Matrix ein hydrophobes Polymer und ein Hydrophobierungsmittel enthält, wobei das Hydrophobierungsmittel aus der Gruppe bestehend aus Cetanol, Cetostearylalkohol, Cetylpalmitat, Wachsen wie Carnaubawachs, Paraffin, Magnesiumstearat, Natriumstearylfumarat und mittel- oder langkettigen Glycerinestern alleine oder Gemischen davon ausgewählt ist.

14. Dosisform nach Anspruch 13, **dadurch gekennzeichnet, daß** das hydrophobe Polymer aus der Gruppe bestehend aus Polyvinylchlorid, Ethylcellulose, Polyvinylacetat und Acrylsäure-Copolymeren ausgewählt ist.

15. Dosisform nach Anspruch 13 oder 14, **dadurch gekennzeichnet, daß** die hydrophobe Matrix ferner eine schwer lösliche oder weniger lösliche Komponente aus der Gruppe bestehend aus Natriumaluminiumsilicat, Calciumphosphat, Aerosil, Titandioxid und Magnesiumcarbonaten enthält.

16. Dosisform nach einem der Ansprüche 1-15, **dadurch gekennzeichnet, daß** es sich bei dem H⁺,K⁺-ATPase-Inhibitor um ein Magnesiumsalz des (-)-Enantiomers von Omeprazol handelt.

17. Dosisform nach einem der Ansprüche 1-16 zur Verwendung bei der Therapie.

18. Verfahren zur Herstellung einer magensaftresistent beschichteten Dosisform gemäß Anspruch 1, die einen H⁺,K⁺-ATPase-Inhibitor und eine hydrophile Matrix oder eine hydrophobe Matrix, die zu einer verlängerten Freigabe des H⁺,K⁺-ATPase-Inhibitors für mindestens 2 und höchstens 12 Stunden führt, und gegebenenfalls pharmazeutisch annehmbaren Hilfsstoff enthält, **dadurch gekennzeichnet, daß** man

a) ein Kernmaterial, das den H⁺,K⁺-ATPase-Inhibitor und die hydrophile Matrix bzw. hydrophobe Matrix und gegebenenfalls pharmazeutisch annehmbaren Hilfsstoff enthält, formt,

b) gegebenenfalls auf das Kernmaterial eine Trennschicht aufbringt und

c) auf das Kernmaterial aus Schritt a) bzw. b) eine magensaftresistente Schicht aufbringt.


19. Verwendung einer oralen pharmazeutischen Zusammensetzung gemäß einem der Ansprüche 1-17 bei der Herstellung eines Arzneimittels mit verbesserter Inhibierung der Magensäuresekretion.

20. Verwendung einer oralen pharmazeutischen Zusammensetzung gemäß einem der Ansprüche 1-17 bei der Herstellung eines Arzneimittels mit verbesserter therapeutischer Wirkung bei der Behandlung von mit übermäßiger Magensäuresekretion einhergehenden gastrointestinalen Störungen.


Revendications

1. Forme de dosage pharmaceutique à libération prolongée et à enrobage gastrorésistant d'un inhibiteur de la H⁺,K⁺-ATPase, **caractérisée en ce que** la forme de dosage comprend une matière de coeur à matrice hydrophile ou


à matrice hydrophobe résultant en une libération prolongée de l'inhibiteur de la H^+, K^+ -ATPase pour un minimum de 2 et un maximum de 12 heures et l'inhibiteur de la H^+, K^+ -ATPase, et en ce que l'inhibiteur de la H^+, K^+ -ATPase est un composé de formule I, un sel alcalin de celui-ci, l'un des énantiomères uniques de celui-ci ou un sel alcalin de l'un des énantiomères d'un composé de formule I




dans laquelle
Het₁ est




ou




Het₂ est




ou



X est



ou



où

N dans le motif benzimidazole signifie que l'un des atomes de carbone du cycle substitués par R₆-R₉ peut être éventuellement remplacé par un atome d'azote dépourvu de substituant ;

R₁, R₂ et R₃ sont identiques ou différents et choisis parmi hydrogène, alkyle, alcoxy éventuellement substitué par fluor, alkylthio, alcoxyalcoxy, dialkylamino, pipéridino, morpholino, halogène, phényle et phénylalkoxy ;

R₄ et R₅ sont identiques ou différents et choisis parmi hydrogène, alkyle et arylalkyle ;

R₆ est hydrogène, halogène, trifluorométhyle, alkyle ou alcoxy ;

R₆-R₉ sont identiques ou différents et choisis parmi hydrogène, alkyle, alcoxy, halogène, halogénoalcoxy, alkyl-carbonyle, alcoxy-carbonyle, oxazolinyle et trifluoroalkyle, ou des groupements adjacents R₆-R₉ forment des structures cycliques qui peuvent être substituées davantage ;

R₁₀ est hydrogène ou forme une chaîne alkylène conjointement avec R₃ ; et

R₁₁ et R₁₂ sont identiques ou différents et choisis parmi hydrogène, halogène ou alkyle.

2. Forme de dosage selon la revendication 1, **caractérisée en ce que** l'inhibiteur de la H^+, K^+ -ATPase est un composé choisi parmi le groupe constitué par l'oméprazole, un sel alcalin de l'oméprazole, l'énantiomère (-) de l'oméprazole et un sel alcalin de l'énantiomère (-) de l'oméprazole.

3. Forme de dosage selon la revendication 2, **caractérisée en ce que** le sel alcalin est un sel de magnésium.

4. Forme de dosage selon la revendication 1, **caractérisée en ce que** l'inhibiteur de la H^+, K^+ -ATPase est un composé choisi parmi le groupe constitué par le lansoprazole, le pantoprazole, des sels alcalins de ceux-ci, un énantiomère unique de ceux-ci et un sel alcalin de ceux-ci.

5. Forme de dosage selon l'une quelconque des revendications 1-4, **caractérisée en ce que** la matière de coeur est recouverte d'une couche de séparation, qui est présente sous la couche d'enrobage gastrorésistant.

6. Forme de dosage selon l'une quelconque des revendications 1-5, **caractérisée en ce que** la matière de coeur comprend en outre des excipients pharmaceutiquement acceptables.

7. Forme de dosage selon l'une quelconque des revendications 1-6, **caractérisée en ce que** la matière de coeur comprend en outre des additifs alcalins.

8. Forme de dosage selon l'une quelconque des revendications 1-7, **caractérisée en ce que** la matière de coeur comprend un germe recouvert de l'inhibiteur de la H^+, K^+ -ATPase et de la matrice hydrophile ou hydrophobe, et éventuellement d'excipients pharmaceutiquement acceptables.

9. Forme de dosage selon l'une quelconque des revendications 1-8, **caractérisée en ce que** la matière de coeur génère un microenvironnement autour de l'inhibiteur de la H^+, K^+ -ATPase non inférieur à $pH = 7$.

10. Forme de dosage selon l'une quelconque des revendications 1-9, **caractérisée en ce que** la matrice hydrophile comprend un polymère hydrophile choisi parmi le groupe constitué par : l'hydroxypropyl méthylcellulose, l'hydroxypropyl cellulose, l'éthylhydroxy éthylcellulose, l'hydroxyéthyl cellulose, la carboxyméthyl cellulose, la carboxyméthyl cellulose de sodium, la méthylcellulose, les oxydes de polyéthylène, la polyvinylpyrrolidone, les alcools polyvinyliques, la gomme adragante et xanthane, ou des mélanges quelconques de ceux-ci.

11. Forme de dosage selon la revendication 10, **caractérisée en ce que** le polymère hydrophile est l'hydroxypropyl méthylcellulose.

12. Forme de dosage selon la revendication 10 ou 11, **caractérisée en ce que** la matrice hydrophile comprend en outre une charge, telle que le silicate de sodium et d'aluminium.

13. Forme de dosage selon l'une quelconque des revendications 1-9, **caractérisée en ce que** la matrice hydrophobe comprend un polymère hydrophobe et un agent d'hydrophobie, l'agent d'hydrophobie étant choisi parmi le groupe constitué par : le cétanol, l'alcool cétostéarylique, le palmitate de cétyle, les cires telles que la cire de carnauba, la paraffine, le stéarate de magnésium, le stéaryl fumarate de sodium et les esters de glycérol à chaîne moyenne ou longue seuls ou des mélanges de ceux-ci.

14. Forme de dosage selon la revendication 13, **caractérisée en ce que** le polymère hydrophobe est choisi parmi le groupe constitué par : le chlorure de polyvinyle, l'éthylcellulose, l'acétate de polyvinyle et les copolymères de l'acide acrylique.

15. Forme de dosage selon les revendications 13 ou 14, **caractérisée en ce que** la matrice hydrophobe comprend en outre un composant peu soluble ou moins soluble choisi parmi le groupe constitué par : le silicate de sodium et d'aluminium, le phosphate de calcium, l'Aerosil, le dioxyde de titane et les carbonates de magnésium.

16. Forme de dosage selon l'une quelconque des revendications 1-15, **caractérisée en ce que** l'inhibiteur de la H^+, K^+ -ATPase est un sel de magnésium de l'énantiomère (-) de l'oméprazole.

17. Forme de dosage selon l'une quelconque des revendications 1-16, pour une utilisation en thérapie.

18. Procédé de fabrication d'une forme de dosage à enrobage gastrorésistant selon la revendication 1, comprenant

un inhibiteur de la H^+, K^+ -ATPase et une matrice hydrophile ou une matrice hydrophobe résultant en une libération prolongée de l'inhibiteur de la H^+, K^+ -ATPase pour un minimum de 2 et un maximum de 12 heures, et éventuellement un excipient pharmaceutiquement acceptable, **caractérisé en ce que** le procédé comprend les étapes suivantes :

a) la mise en forme d'une matière de coeur, comprenant l'inhibiteur de la H^+, K^+ -ATPase et la matrice hydrophile ou la matrice hydrophobe, et éventuellement des excipients pharmaceutiquement acceptables,

b) éventuellement, l'application d'une couche de séparation sur la matière de coeur, et

c) l'application d'une couche d'enrobage gastrorésistant sur la matière de coeur de l'étape a) ou de l'étape b).

19. Utilisation d'une composition pharmaceutique orale selon l'une quelconque des revendications 1-17, dans la fabrication d'un médicament présentant une inhibition améliorée de la sécrétion d'acide gastrique.

20. Utilisation d'une composition pharmaceutique orale selon l'une quelconque des revendications 1-17, dans la fabrication d'un médicament présentant un effet thérapeutique amélioré dans le traitement de troubles gastro-intestinaux associés à une sécrétion excessive d'acide.